


# Progress on Application of Modal Aerosol Dynamics to CAM

Xiaohong Liu, Steve Ghan, Richard Easter, Rahul Zaveri, Yun Qian  
(*Pacific Northwest National Laboratory*)

Jean-Francois Lamarque, Peter Hess, Natalie Mahowald, Francis Vitt,  
(*NCAR*)



# Current Aerosol Treatment in CAM3

sulfate

hydrophobic  
black  
carbon

sea salt 1

soil dust 1

ammonium

hydrophobic  
organic  
carbon

sea salt 2

soil dust 2

nitrate

hydrophilic  
black  
carbon

sea salt 3

soil dust 3

secondary  
organic  
carbon

hydrophilic  
organic  
carbon

sea salt 4

soil dust 4

## Current Weaknesses in CAM

- Aerosol species are externally mixed (individual particles are composed of only a single species).


# Current Weaknesses in CAM

- Aerosol species are externally mixed (individual particles are composed of only a single species).
- Their size distribution is prescribed (number is diagnosed from the predicted mass).
  - Processes that should only affect mass (condensation, chemistry) also affect number.
  - Processes that only affect number (nucleation, coagulation) are neglected.

# Current Weaknesses in CAM

- Aerosol species are externally mixed (individual particles are composed of only a single species).
- Their size distribution is prescribed (number is diagnosed from the predicted mass).
  - Processes that should only affect mass (condensation, chemistry) also affect number.
  - Processes that only affect number (nucleation, coagulation) are neglected.
- Hydrophobic carbon ages to hydrophilic with prescribed timescale

# Proposed Benchmark Aerosol Treatment for CAM4




# Proposed Simplified Aerosol Treatment for CAM4

Assume aerosol are hydrated for  $RH >$  crystallization  $RH$ .

Carry soil dust and sea salt in same mode because sources are separate.

Assume primary carbon is internally mixed with secondary aerosol.

Assume ammonium neutralizes sulfate, and neglect nitrate.



All modes are log-normal with prescribed width.

Total transported aerosol tracers: 15

Cloud-borne aerosol are predicted but not transported.

# Progress

- Emissions (size-resolved) (done)
  - ✓ Fossil fuel SO<sub>2</sub> emissions over 0-100 m and >100 m based on EDGAR-2000;
  - ✓ 3% (by mole) of fossil fuel SO<sub>2</sub> emissions goes to SO<sub>4</sub> (Aitken & Accu. modes, mass & number, Whitby 1978);
  - ✓ OM and BC to primary carbon mode (mass & no.);
  - ✓ Sea salt emission based on Gong et al (1997) to fine and coarse sea salt modes (mass & no.);
  - ✓ Dust emissions to fine and coarse dust modes (mass & no.).

# Progress

- Aerosol activation and droplet nucleation (Ghan scheme) (done)
- Wet scavenging (done)
  - ✓ In-cloud rainout based on activated aerosol;
  - ✓ Below-cloud impaction scavenging rates (mass & no.) using a look-up table (wet size, precipitation rate).
- Dry deposition (based on Zhang et al., 2001) & gravitational settling (done)
- Cloud sulfur chemistry (done)
  - ✓ Bulk sulfate chemistry based on calculated pH
  - ✓ Sulfate mass produced distributed to modes based on number of activated aerosols in modes.
  - ✓ Include contribution from  $\text{H}_2\text{SO}_4$  (g) uptakes

## Progress

- New particle formation: binary nucleation ( $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$ ) parameterization (Vehkamäki et al., 2002) **(done)**
- Coagulation: Brownian within, between modes. **(done)**
- Water vapor and trace gas ( $\text{H}_2\text{SO}_4$  &  $\text{NH}_3$ ) uptake **(done)**
- Intermode transfer (renaming) due to condensation, coagulation, and cloud chemistry **(done)**
- **Compile OK**

## Works in Progress

- Trace gas and water vapor uptake (with hysteresis dependent on previous aerosol water) will be replaced with MOSAIC module (consider gas-particle transfer)
- Aerosol optical properties: parameterization in terms of wet refractive index and wet surface mode radius.

# CAM Simulations

- Benchmark modal present-day
  - On-line oxidants ( $\text{HO}_x, \text{O}_3$  chemistry)
  - Off-line oxidants (input  $\text{HO}_x, \text{O}_3$ )
- Benchmark modal pre-industrial
- Simplified modal present day, offline oxidants
- Simplified modal pre-indu., offline oxidants
- Offline benchmark present day (interpolation from monthly mean of on-line benchmark)
- Offline benchmark pre-industrial

# Testing

- Evaluate on-line benchmark treatment using *in situ* (surface and aloft) and remote aerosol measurements (mass, number, size, CCN, AOD, ...). Utilize AEROCOM and other model evaluation efforts.
- Evaluate approximations used in on-line simplified treatment by comparing direct and indirect aerosol effects with on-line benchmark treatment.
- Evaluate off-line benchmark treatment by comparing with on-line benchmark treatment.