



Community Earth
System Model

**Be
Respectful**

**Be
Collaborative**

**Consider
differing
opinions**

**Communicate
openly**

**Be
mindful
of others**



NCAR
OPERATED BY UCAR

Community Earth
System Model

CESM Communications:

<https://www.cesm.ucar.edu/communications>

Subscribe to the CESM Newsletter :

www.cesm.ucar.edu/news/newsletters



@ncar_cgd






@NCAR_CGD



NSF NCAR CGD



@ncar-cgd.bsky.social




Updates on ocean biogeochemistry in CESM3

Kristen Krumhardt
NSF NCAR
BGCWG co-chair

Contributions from Keith Lindsay, Mike Levy, Nicola Wiseman,
Keith Moore, Jun Yu, Gustavo Marques, Ian Grooms



The background of the slide is a grid of 20 oceanographic cross-sections. Each section shows a vertical profile of the ocean with different colors representing various water masses, biological features, or physical properties. The colors range from deep blues and purples to bright yellows and oranges. The sections are arranged in a 4x5 grid, with the text overlaid on the left side.


Overview of CESM3 MARBL developments

- New planktonic ecosystem
- Coccolithophores
- Fully variable phytoplankton stoichiometry
- New iron cycle updates
- Fixing OMZ problem


4 Phytoplankton / 2 Zooplankton ("4p2z") MARBL

- CESM2 had 3p1z plankton
- Small phytoplankton and diatoms do most ocean NPP
- Coccolithophores and diazotrophs have important BGC roles
- 2 zooplankton types allows offline coupling to a fish model


CESM-MARBL-4p2z planktonic ecosystem




4p2z ecosystem allows offline coupling to fish model



Coccolithophore phytoplankton functional type & ocean acidification




- Coccolithophore importance to carbon cycle
 - Affect air-sea CO₂ uptake via calcification
 - CaCO₃ shell ballasts organic matter to deep ocean
- Ocean acidification impacts coccolithophore calcification and growth




Norway

As more CO₂ absorbs into the ocean, cocco calcification decreases



$$\text{CaCO}_3 \rightarrow \frac{\text{Particulate Inorganic Carbon (PIC) shell}}{\text{Particulate Organic Carbon (POC) inside}} = \frac{\text{PIC}}{\text{POC}}$$





- Based on a large meta-analysis of laboratory experiments (Krumhardt et al., 2017, *Progress in Oceanography*)
- Less calcification → more alkalinity stays in the surface ocean → more CO₂ uptake/less ballasting of carbon to deep ocean

Fully variable C:N:P:Fe:Si stoichiometry in phytoplankton


- Variable phytoplankton N:C new in CESM3
- Each PFT has min/max of X:C

Small phyto N:C in MOM6-MARBL development run



Fully variable C:N:P:Fe:Si stoichiometry in phytoplankton

- Variable phytoplankton N:C new in CESM3
- Each PFT has min/max of X:C
- Improved match to particulate organic matter stoichiometry →
- Strong implications for ocean biological carbon pump to climate change





Iron cycle updates

(Keith Moore & Jun Yu, UCI)

- Iron is an important control on phytoplankton growth in many regions



Diatom most limiting nutrient in MOM6-MARBL development run



Iron cycle updates

(Keith Moore & Jun Yu, UCI)



- Iron is an important control on phytoplankton growth in many regions
- Iron comes from many sources, updated in CESM3





Iron cycle updates

(Keith Moore & Jun Yu, UCI)

- Iron is an important control on phytoplankton growth in many regions
- Iron comes from many sources, updated in CESM3
- For atmospheric deposition, Coarse/Fine ratio determines soluble %
- Scavenging of iron in the ocean also updated for CESM3



Leith-E backscatter helps equatorial OMZs




Backscatter generates more realistic zonal jets: stacked at the equator, with Tsuchiya jets on the flanks

Effect of using Leithy backscatter on OMZs


O_2 at 500m

20 $mmol\ m^{-3}\ O_2$
contour



CESM Control




Observations



CESM with backscatter





CESM Control O_2 bias




Overly extensive
OMZ looks better

CESM Backscatter O_2 bias




Leith-E backscatter helps equatorial O₂ and NO₃ biases

O₂



145: no backscatter
152: backscatter

NO₃





- Low O₂ triggers denitrification - remineralization of POC that consumes NO₃

Future plans for subsequent versions of CESM3

- 8 phytoplankton / 4 zooplankton MARBL

- Ocean biogeochemistry coupling with sea ice



Thank you! Any questions?



Email: kristenk@ucar.edu



NCAR
OPERATED BY UCAR