

E3SMv3: Wave Configuration Current Status + Future Plans

Erin Thomas

Theoretical Division  Los Alamos National Laboratory

*Andrew Roberts, Elizabeth Hunke,
Olawale Ikuyajolu, Steven Brus, Luke Van Roekel,
Jon Wolfe, Adrian Turner
+ CESM Wave+Ice Folks*




PCWG meeting. NCAR, Boulder, CO.
5-6 Feb 2026

LAUR 25-26182



- **E3SMv3: Overview**
- **E3SMv3: Recent (Relevant) Developments**
 - WAVEWATCHIII Unstructured Mesh
 - Wave – Ocean – Atm Momentum Coupling
 - Wave – Sea Ice Coupling
- ***E3SMv3: Ongoing Efforts + Future Plans***
 - Resolution
 - AI / ML for Waves
 - other big picture ideas for E3SM

- **E3SMv3: Overview**
- **E3SMv3: Recent (Relevant) Developments**
 - WAVEWATCHIII Unstructured Mesh
 - Wave – Ocean – Atm Momentum Coupling
 - Wave – Sea Ice Coupling
- ***E3SMv3: Ongoing Efforts + Future Plans***
 - Resolution
 - AI / ML for Waves
 - other big picture ideas for E3SM



Unstructured ocean-sea ice (blue) and waves (black) meshes



Flagship Earth system model developed across DOE labs



Now includes ice shelves, sea ice floes and wave attenuation



Focused on seasonal to multi decadal prediction



Targeting exascale computing using GPUs & AI




E3SMv3 (2024) 30km/100km/0.5° marine/atmos/land



E3SMv4 (2027) eddy & cloud resolving target



Current Capabilities in the Energy Exascale Earth System Model (E3SMv3):

- E3SM is a *fully coupled* global Earth system model targeting exascale computing
- **Now Includes:**
 - **Fully coupled Waves (WW3)**
 - **Wave-Ice Interactions (FSD, Icepack)**
 - Antarctic ice shelves
- **Unstructured meshes: capable of regional refinement in the sea ice, ocean, and waves**




- **E3SMv3: Overview**
- **E3SMv3: Recent (Relevant) Developments**
 - WAVEWATCHIII Unstructured Mesh
 - Wave – Ocean – Atm Momentum Coupling
 - Wave – Sea Ice Coupling
- *E3SMv3: Ongoing Efforts + Future Plans*
 - Resolution
 - AI / ML for Waves
 - other big picture ideas for E3SM

Resolution = 30 km along coastlines
to 200+ km in open ocean




Unstructured ocean-sea ice (blue)
and wave (black) meshes

Introduction



Wave - Atmosphere Coupling



Neutral drag coefficient (C_{DN})

NCAR	$(2.7U_{10N}^{-1} + 0.142 + 0.0.0764U_{10N}) \times 10^{-3}$
COR3.0a	$C_{DN}[z_{o,m}[\alpha(U_{10N})]]$
COR3.0a-WAV	$C_{DN}[z_{o,m}(\underline{\alpha}_w)]$

E3SM default setting = NCAR

No possible wave feedback to the atm since the *drag coefficient is parametrized as a function of only 10m wind speed.*

	Neutral drag coefficient (C_{DN})
NCAR	$(2.7U_{10N}^{-1} + 0.142 + 0.00764U_{10N}) \times 10^{-3}$
COR3.0a	$C_{DN}\{z_{o,m}[\alpha(U_{10N})]\}$
COR3.0a-WAV	$C_{DN}\{z_{o,m}(\underline{\alpha}_w)\}$


The COARE 3.0 turbulent flux scheme **would** allow for wave coupling through the Charnock parameter

-HOWEVER-


- 1) this flux scheme is **NOT** used in standard E3SM simulations, so any results are NOT comparable to baseline E3SM simulations.
- 2) The flux scheme over sea ice is based on the NCAR, so the fluxes over ocean and ice surfaces would be incompatible


	Neutral drag coefficient (C_{DN})
NCAR	$(2.7U_{10N}^{-1} + 0.142 + 0.0.0764U_{10N}) \times 10^{-3}$
COR3.0a	$C_{DN}\{z_{o,m}[\alpha(U_{10N})]\}$
COR3.0a-WAV	$C_{DN}\{z_{o,m}(\alpha_w)\}$

Therefore, we created a 'modified NCAR' surface flux scheme that allows for wave coupling by **calculating the surface drag coefficient from the surface roughness (Z0) directly from WW3:**



Wave - Ocean Coupling






- Partitioned Stokes drift values computed in WW3
- Partitioned Stokes drift passed from WW3 to MPAS-Ocean
- the full depth profile of the Stokes drift then reconstructed in the ocean model
- In MPAS-O, reconstructed Stokes drift used to estimate the Langmuir enhancement factor.

Figure from Rühls, S., et. al., Ocean Sci., 2025


Default Atm-Ocean coupling setup assumes ocean side stress is equal to atm stress....

A.



$$\tau_{ocn} = \tau_{atm}$$

B.




However: some of Atm momentum is lost to waves ... and waves themselves can impart a stress to the ocean... **We now account for all of this**

A.




$$\tau_{ocn} = \tau_{atm}$$

B.



Wave - Sea Ice Coupling




Sea Ice Floe Size Distribution (FSD)

- ICEPACK column physics

$$\frac{\partial f(r, h)}{\partial t} = - \overbrace{\nabla \cdot (f(r, h) \mathbf{v})}^{\text{Advection}} + \underbrace{\mathcal{L}_T}_{\text{Thermodynamics}} + \underbrace{\mathcal{L}_M}_{\text{Mechanical}} + \underbrace{\mathcal{L}_W}_{\text{Waves}}$$

Floe Breaking Due to Waves

- Default FSD breaking scheme in Icepack
(Horvat and Tziperman 2015/Roach et al. 2018, 2019)
- Sea ice flexes with the sea surface height field
- Ice Floe breaks if 'strain' exceeds critical threshold



Wave Attenuation due to Sea Ice: Meylan et al 2021

- Dissipation of wave power due to sea ice
 - Low Frequency waves (long wavelengths) dissipate LESS in sea ice (and are critical to wave-ice coupling in models)





Figure 7
Meylan et al. 2021


50-Yr Annual Mean MLD

MLD (ANN, years 0001-0050)

WaveMomentum_PIControl_WW3v6.07



Control: WaveMomentum_PIControl_NoWave




Wave
Ocean
Atm
Coupling

No
Wave

**Wave-Ocean-Atm coupling:
MLD increases
(Stokes/ Langmuir)**


Main - Control



50-Yr Annual Mean SST


SST (ANN, years 0001-0050)

WaveMomentum_PIControl_WW3v6.07



Wave
Ocean
Atm
Coupling


Control: WaveMomentum_PIControl_NoWave



No
Wave



**Wave-Ocean-Atm coupling:
increased mixing = cooling**

Main - Control




**50-Yr JFM
Mean Sea Ice
Area**

Wave
Ocean
Atm
Coupling



No
Wave




**Wave-Ocean-Atm
coupling:**

**Cooling = Increased
sea ice area (and
increased thickness)**

**Wave-Ocean-Atm
minus
NoWave**

Main - Control
150°W 180°W 150°E




**50-Yr JFM
Mean Sea Ice Area**

**Wave-Ice
coupling
generally
counteracts
the impacts of
wave-ocean
coupling
(lateral melt)**


**Wave-Ocean-Atm-Ice
minus
Wave-Ocean-Atm**

Main - Control
150°W 180°W 150°E



**Wave-Ocean-Atm
minus
NoWave**

Main - Control




**50-Yr JJA
Mean Sea Ice Area**

**Wave-ice
coupling
generally
counteracts
the impacts of
wave-ocean
coupling
(lateral melt)**

**Wave-Ocean-Atm-Ice
minus
Wave-Ocean-Atm**

Main - Control



E3SM: Overview

E3SM: Recent Developments


- WAVEWATCHIII Unstructured Mesh
- Wave- Ocean- Atm Momentum Coupling
- Wave – Sea Ice Coupling

E3SM: Ongoing Efforts + Future Plans


- Resolution
- AI / ML
- S2S / S2D

Ocean
+
Sea Ice

E3SMv3 Standard Resolution



E3SMv3 High Resolution




Standard (left) versus high-resolution (right) E3SM meshes (17-fold data volume difference for surface fields)


E3SM: Goals for Resolution ... Everywhere

Atmosphere: 3km cloud resolving atmosphere

Himawari Satellite Visible Image (Jan 22 2 UTC)



Shortwave Cloud Radiative Effect (Jan 22 2UTC)



Cold-air outbreak off Siberia on January 22, 2020 at 2:00:00 UTC (~local noon) from a Himawari visible satellite image (left) and shortwave cloud radiative effect from SCREAM (right) from convection permitting model by Caldwell et al. (2021)

E3SM: Overview


E3SM: Recent Developments


- WAVEWATCHIII Unstructured Mesh
- Wave- Ocean- Atm Momentum Coupling
- Wave – Sea Ice Coupling

E3SM: Ongoing Efforts + Future Plans

- Resolution
- AI / ML (example for Waves)

WW3 is a **significant** computational cost in LR E3SMv3





- Source Terms accounts for nearly 80% of total compute time
- Dissipation source term ~44% of total
- Non-linear source term ~20% of total

Figure 4. The Callgraph obtained by profiling WAVEWATCHIII with 8 CPUs. Each box includes the name of the subroutine and its relative execution time as a percentage
Ikuyajolu et al. Geosci Model Dev. 2023

Machine Learning Emulation of Non-linear Wave Interactions

Exact (*WRT*) nonlinear wave interactions (S_{nl}) is $10^2 - 10^3$ more costly than Discrete Interaction Approximation (*DIA*) method due to the 6D Boltzmann integral.


- *DIA* sacrifices accuracy for speed

Can we develop a neural network emulator that has the accuracy of the **WRT** and the speed of **DIA** ?

□ ML model


- 4 hidden layers with 3072 neurons
- Trained on wave-spectra (50 freq x 36 dir)
- stand-alone WW3 simulation for 2010

Randomly selected training data locations shown as boxed regions over areas of high and low wave activity.




Significant
Wave
Height



RMSE: APPROX - EXACT




RMSE: ML emulator - EXACT



Peak
Wave
Period







- ***E3SMv3 Wave – Enabled Configuration***
- ***Ongoing Efforts + Future Plans***
Resolution
AI/ML
Shorter time scales (S2S, S2D)

Thank You!